

Conservative surgeries do yield good functional results if properly planned and executed. Such surgeries yield better results in the presence of delayed presentation with complications like infection, callus formation, deformity. Revisiting and doing subsequent elective surgery becomes easy in the absence of infection.

Achieving a good functional result is more rewarding than restoring anatomy and cosmesis at the cost of function, especially in complicated delayed case presentations.

Clinical Message

References

1. Tiwari A, Kanojia R K, Kapoor S K. Surgical management for late presentation of supracondylar humeral fracture in children. *Journal of Orthopaedic surgery*. Aug.1, 2007.
2. H.De Boeck, P.De Smet Valgus Deformity following Supracondylar Elbow fractures in Children *Acta Orthopaedica Belgica*, Vol.63-4-1997.
3. Ating'a, J.E.O.: Conservative Management of Supracondylar Fractures of the Humerus in Eastern Provincial General Hospital, Machokos. *East Afr.Med.J.*, 61:557-560, 1984.
4. Charles A. Rockwood, Jr., M.D., Kaye E.Wilkins, M.D., Richard E.King, M.D. *Textbook of Fractures in Children*, 3rd edition, 588-589, 1991.
5. C. G. Attenborough Remodelling of the Humerus after supracondylar fractures in childhood. *The Journal of Bone and Joint Surgery*, Vol.35B, No.3, August 1953.

Conflict of Interest: Nil
Source of Support: None

How to Cite this Article

Deshpande M M, Punit A S, Rohit V. Management of seventeen day old infected type 4b-Supracondylar fracture Humerus-A Case Report. *International Journal of Surgical Cases* 2015 May-July;1(1): 10-12.